

(i.e., nivolumab). Fourth, significant declines in serum hepatitis B surface antigen (HBsAg) are demonstrated during gene therapy, which have never been seen using the most potent polymerase inhibitors (i.e., tenofovir or entecavir). Finally, unanticipated significant reductions in cccDNA are seen with HBV gene therapy, most likely as prove of an indirect benefit of waning the immunosuppressive effect of large over amounts of HBsAg released by infected hepatocytes that contributes to T-cell exhaustion.

In a pioneering study, Roche was the first to publish the potent effect of an oral small molecule that blocked HBV gene expression (*Mueller et al. J Hepatol 2018;68:412-20*). The drug belonged to the dihydroquinolizinone class, and directly or indirectly modified viral RNAs, promoting their degradation. This post-transcriptional silencing was accompanied by rapid drops in HBV-DNA and more importantly in serum HBsAg in the humanized mice. However, Roche decided to discontinue any further clinical development of the drug.

Nowadays, two major groups of agents are being developed as HBV gene therapies. At this time, interference RNA (iRNA) molecules and nucleic acid polymers (NAPs) are the most promising. Overall, iRNA is double-stranded RNA molecules, 20 nucleotides long. One strand matches a segment of specific HBV mRNA and induces its degradation. Several iRNA molecules have entered into Phase II clinical trials (*Flisiak et al. Exp Op Biol Ther, in press*), including ARB-1467 and AB-729 (Arbutus), ARO-HBV (Arrowhead), ALN-HBV (Alnylam), and IONIS-HBVRx (Ionis). In most cases, they are tested as part of combination therapy with nucleos(t)ide analogs and/or peginterferon.

NAPs are phosphorothioate 40 length oligonucleotides that do not map any HBV sequence. However, they interact with a liver host target protein (apolipoprotein-like) and result in specific inhibition of HBV mRNAs. This is followed by rapid suppression of HBsAg release (*Roehl et al. Mol Ther Nuc Acids 2017;8:1-12*). In a pilot study with intravenous REP-2139, investigators from Replicor demonstrated strong reductions in HBV-DNA along with significant drops in HBsAg and seroconversion in some patients. More interestingly was the recognition

of significant reductions in hepatic cccDNA, most likely a result of an indirect effect following the removal of large amounts of HBsAg from the bloodstream that contributes to impaired T-cell responses in chronic hepatitis B patients (*Bazinet et al. EASL, Paris 2018; abstract FRI-343*). An improved NAP, named REP-2165 and subcutaneous administration are currently being tested.

Vicente Soriano

*La Paz University Hospital and UNIR Health Sciences School,
Madrid, Spain*

Dr. Robert Redfield Appointed as New CDC Director

Well-known AIDS researcher Robert Redfield picked to lead the CDC on March 2018. He is one of the HIV/AIDS pioneers in the United States. During two decades at the Walter Reed Army Institute of Research, in Bethesda, MD, he made pivotal contributions, highlighting the importance of heterosexual HIV transmission, developing the Walter Reed staging system for HIV infection, and demonstrating that active HIV replication takes place during all stages of HIV disease.

In 1996, he was one of the cofounders of the Institute of Human Virology in Baltimore, MD. Nowadays, he was running a treatment network for HIV and hepatitis C patients. With this background, he would be well prepared to combat one of the DHHS and CDC's top priorities, the opioid epidemic (*Kolodny et al. JAMA 2017;318:1537-8*).

In an interview, Robert Gallo, director of the Institute of Human Virology, said "Redfield is a dedicated and compassionate physician who truly cares about his patients and is deeply committed to ensuring patients receive the highest quality of care possible. Dr. Redfield has served his country well, and consistently demonstrates strong public health instincts that are grounded in science and clinical medicine. In my view, despite the loss to our institute, I believe this makes him the ideal candidate to direct the CDC".

Pablo Barreiro
*La Paz University Hospital
Madrid, Spain*